

POLICY: 555.21
 TITLE: Pediatric Airway Obstruction by Foreign Body
 EFFECTIVE: 02/01/2026
 REVIEW: 02/2028
 SUPERCEDES:

APPROVAL SIGNATURES ON FILE IN EMS OFFICE

PAGE: 1 of 2

PEDIATRIC AIRWAY OBSTRUCTION BY FOREIGN BODY

- I. AUTHORITY
Health and Safety Code, Division 2.5, California Code of Regulations, Title 22, Division 9
- II. PURPOSE
To serve as a patient treatment standard for EMRs, EMTs, and Paramedics within their scope of practice.
- III. PROTOCOL
If there is a history of a febrile illness and copious drooling, strongly consider epiglottitis. If epiglottitis is suspected and patient is ventilating adequately, transport immediately and avoid visualization of the airway.

**Provider Key: F = First Responder/EMR E = EMT O = EMT Local Optional SOP
 P = Paramedic D = Base Hospital Physician Order Required**

	F	E	O	P	D
ASSESSMENT: look for signs of poor perfusion or respiratory distress (delayed capillary refill, diminished distal pulses, cool extremities, ALOC).	X	X	X	X	
OXYGEN: 100% by non-rebreather mask or blow-by.	X	X	X	X	
PULSE OXIMETRY: apply and monitor.		X	X	X	
ECG MONITOR: lead placement may be delegated. Treat as indicated.				X	
CONSCIOUS PATIENT - ABLE TO SPEAK, COUGH OR CRY					
REASSURE PATIENT: encourage coughing.	X	X	X	X	
SUCTION: as needed to control secretions.	X	X	X	X	
CONSCIOUS PATIENT - UNABLE TO SPEAK, COUGH, OR CRY					
BACK BLOWS & CHEST THRUSTS: for patients < 1 year of age. Alternate back blows and chest thrusts with head inferior to chest. For patients >1 year of age, use abdominal thrusts in conjunction with back blows.	X	X	X	X	
REASSESS: repeat basic airway maneuvers until obstruction is cleared or the patient becomes unconscious.	X	X	X	X	
UNCONSCIOUS PATIENT					
VISUALIZE AIRWAY: use laryngoscope and pediatric Magill Forceps. May finger sweep only if obstruction visible. Reassess airway prior to ventilation after each CPR cycle.				X	
CHEST THRUSTS/HP-CPR: 15:2 ratio, even if pulses are present.	X	X	X	X	
BLS AIRWAY: okay if airway patent. Support ventilations with appropriate airway adjuncts.	X	X	X	X	
SUPRAGLOTTIC AIRWAY: if GCS is < 8 and not rapidly improving.			X	X	
CAPNOGRAPHY: apply and monitor if SGA has been placed.				X	

	F	E	O	P	D
NEEDLE CRICOTHYROTOMY: if unable to ventilate with SGA <ul style="list-style-type: none"> ● Quicktrach Child device for patients 10-35 kg (22-77 lbs). ● Quicktrach device for patients > 35 kg (> 77 lbs). ● 14 – 18G catheter for patients < 10kg (< 22 lbs). ● Ventilate with high flow oxygen. 				X	

Refer to 555.23 PEDIATRIC RESPIRATORY DISTRESS.

NOTE: Transport patient immediately to the closest receiving hospital if unable to clear obstruction or otherwise establish an airway. All patients should be transported to a receiving hospital regardless of airway maneuvers.

Pediatric Normal Vital Signs

Age	HR	RR	BP	Temp (C)	Temp (F)
<i>Premie</i>	120-170	40-70	55-75/35-45	36-38	96.8-100.4
<i>0-3 months</i>	100-160	35-60	65-85/45-55	36-38	96.8-100.4
<i>3-6 months</i>	90-120	30-45	70-90/50-65	36-38	96.8-100.4
<i>6-12 months</i>	80-120	25-40	80-100/55-65	36-38	96.8-100.4
<i>1-3 years</i>	70-110	20-30	90-105/55-70	36-38	96.8-100.4
<i>3-6 years</i>	65-110	20-25	90-110/60-75	36-38	96.8-100.4
<i>6-12 years</i>	65-100	14-22	90-120/60-75	36-38	96.8-100.4
<i>12+</i>	55-100	12-20	100-135/65-85	36-38	96.8-100.4